

Glucan and Humic Acid: Synergistic Effects on the Immune System

Vaclav Vetvicka, Roberto Biggori,² Angelo M. Zamaretti,²
Jose M. Garcia-Mina,^{2,3} and Jean-Claude Viret⁴

¹Department of Pathology, University of Louisville, Louisville, Kentucky, USA; ²R&D Department, Timac Agro International, Breda, Italy; ³Department of Chemistry and Soil Chemistry, University of Navarra, Pamplona, Spain; and ⁴INRA, Department of Animal Agronomy, Jouy en Josas, France

ABSTRACT Humic acids are natural compounds resulting from decomposition of organic matter. In spite of their omnipresence, our knowledge of their biological effects is limited, and current findings are controversial. We decided to evaluate the immunological effects of two different types of humic acids, differing in source and physicochemical characteristics, along with both components either alone or in combination with the well-established yeast-derived immunomodulator glucan (as measured by their effects on both the cellular (phagocytosis and tumour necrosis factor) and humoral (antibody production and cytokine secretion) branches of the immune system). In summary, our results suggest that humic acids are biologically active immunomodulators affecting both the humoral and cellular branches of immune reactions. In addition, the two humic acids studied here are working in synergy in stimulation of the immune reaction, supporting further studies of these natural immunomodulators.

KEY WORDS: glucan, humic acids, immunity, phagocytosis

INTRODUCTION

The immunomodulatory activity of glucan has been well documented for over 50 years. The first interest in the immunomodulatory properties of polysaccharides arose after experiments showing that a sterile yeast cell preparation stimulated macrophages via activation of the complement system.¹ Further work identified the active component as β -D-glucan.² Numerous studies have subsequently shown that β -D-glucans, given in particulate or soluble form, have both immunostimulating properties, such as inducing granulocyte and dendritic cell activation.^{3,4} More than 2,500 publications have reported that β -D-glucan can either solely or in combination with other immunomodulatory ingredients. Currently, glucans are considered to be one of the most efficient biological immunomodulators (for review, see Novak and Vetvicka⁵).

Some studies suggested that glucan molecules have synergistic effects when combined with glucan. Numerous reports have shown some beneficial effects when glucan was given in combination with vitamins C and E. The main reason why vitamins C and E synergize is the fact that this combination has been shown to stimulate the expression of immune responses as well as glucan, in terms of phagocytic activities, natural

killer cell activity, and specific antibody formation. A mouse study showed synergistic effects of glucan and vitamin C combination in the treatment of infection by *Mesocricetus* *steieri*; the combination resulted in a positive modulation of the fibroblast and physiological changes.⁶ The same group found previously that a yeast-derived glucan preparation was effective against several helminth parasites.⁷ With respect to the liver, glucan is known to help against ischemic reperfusion injury of the liver.⁸ The mechanisms of these effects are probably due to the glucan-induced decrease of the expression of immediate early genes following reperfusion.⁸

Humic substances occur mainly in heavily degraded peat and humic acids (HAs) are present in a wide range of high-molecular-weight macromolecular substances, such as complex polysaccharides, lignin, and tannins. Together with fatty acids, they represent a certain fraction of the group of organic compounds called humic substances, which are sometimes considered to be toxic.⁹ More detailed studies revealed a controversial result: high doses of HA induced chromosomal abnormalities in lymphocytes,¹⁰ whereas a low dose of HA stimulated humoral and cellular immune responses.¹¹ On the other hand, several immunostimulating potentialities have been reported for biological activities, were recently established in various types of HAs, including their antitumor properties, and proliferation of lymphocytes.¹⁶ Also, addition of HA into the feed of experimental animals results in improved growth and health.¹⁷

Manuscript received 29 July 2009; Revision accepted 25 August 2009.

Address correspondence to Vaclav Vetvicka, Department of Pathology and Laboratory Medicine, University of Louisville, 541 South Fourth, MEDR Building, Room 224, Louisville, KY 40202, USA, E-mail: vaclav@iglou.com

The present observations showing that glucan's biological activities were significantly improved by resveratrol led us to evaluate the possible synergistic effects of glucan and HAs on immune reactions.

MATERIALS AND METHODS

Animals

Female 6- to 8-week-old BALB/c mice were purchased from the Jackson Laboratory (Bar Harbor, ME, USA). All animal work was done according to the protocol of the Institutional Animal Care and Use Committee of the University of Louisville, Louisville, KY, USA. Animals were sacrificed by CO₂ asphyxiation.

Materials

RPMI 1640 medium, sodium citrate, bovine hemin, antibiotics, Wright's stain, *Ultraspeed* test kit, FETOXADT Freund's adjuvant, and concanavalin A (ConA) were obtained from Sigma Chemical Co. (St. Louis, MO, USA). Fetal calf serum was from Hyclone Laboratories (Logan, UT, USA).

β -D-Glucans

The glucans used in this study were purchased from the following companies: yeast-derived β -D-glucan (Aurum) from 300z from Transfer Point (Goldschmidt, USA) and soluble glucan laminarin from Sigma.

Extraction procedure

The HAs were used in the experiments were extracted from denardite obtained from Czech Republic (HZ) and antileonite extracted from lignite obtained from China (HLS). The different HAs were isolated, purified following the International Humic Substances Society procedure,^{1,2} and freeze-dried.

Solid-state ¹³C nuclear magnetic resonance (NMR) spectroscopy

Solid-state ¹³C NMR spectra were obtained on a Bruker (Billerica, MA, USA) Avance 400WB (9.4 T) spectrometer at 100.47 MHz using the cross-polarization magic angle spinning technique with a spinning speed of 12 kHz, 90° pulse width of 10 microseconds, acquisition time of 4.0 second delay.

Elemental analysis

The carbon, hydrogen, and nitrogen contents of the lyophilized samples were analyzed in duplicate by a LECO® (St. Joseph, MO, USA) COHN 200 analyzer. The oxygen content was determined by difference (ash-free basis).

Phagocytosis in vitro

Phagocytosis was measured *in vitro* using synthetic microspheres (2-hydroxyethyl methacrylate [HEMA] parti-

cles) after intraperitoneal injection of glucan and/or HAs described in the Introduction. The cells for cell adhesion were incubated with 0.01 ml of HEMA particles (5 × 10⁸ particles). The cells were incubated at 37°C for 60 minutes with intermittent shaking. Smears were stained with Wright's stain. Cells with three or more HEMA particles were considered positive. The same smears were also used for evaluation of cell types.

Evaluation of interleukin (IL)-2 production

Purified spleen cells (2 × 10⁶ cells in RPMI 1640 medium with 5% fetal calf serum) were added into wells of 24-well tissue culture plate. After addition of 10⁶ ConA into positive control wells, cells were incubated for 72 hours in a humidified incubator at 37°C and 5% CO₂. At the end point of the incubation, supernatants were collected, filtered (pore size 0.45 μm) and tested for the presence of IL-2. IL-2 levels were measured using a Quantikine® Mouse IL-2 kit (R&D Systems, Minneapolis, MN, USA).

Cytokine array

Endothelial cytokines were measured in mouse serum by the direct dot blot (Eugene, OR, USA) or mice were injected with the test combination for 24 hours. The mice were sacrificed, and serum was collected and stored in a -80°C freezer. For the cytokine analysis, we used the protein microarray service provided by Adipos Biotech. In brief, the service used a multiplexed antibody-based protein detection multiplex assay. A streptavidin-Cy5-lipid conjugate was used for mass cytometry. The assay was done in quadruplicate by positive and negative control spotted on each microarray. In the assay context, the following cytokines: IL-2, interferon- γ , interferon- α , interferon- β , IL-4, IL-5, interferon-inducible protein-10, macrophage inflammatory protein-1 β , IL-13, IL-1 β , and monocyte chemoattractant protein (MCP).

Tumor inhibition in vivo

Mice were injected directly into the mammary fat pads with 10⁶ of Ptas64 cells in phosphate-buffered saline (PBS) per mouse. The experimental treatment was begun after palpable tumors were found (usually 4–6 days after injection of cells) and after mice were assigned to experimental groups. Experimental treatment was achieved by intraperitoneal injections of diluted samples diluted in PBS. After treatment, the mice were sacrificed, and tumors were removed and weighed.

Antibody formation

Mice were injected twice (2 weeks apart) with 100 μg of ovalbumin and serum was collected 7 days after the last injection. Levels of specific antibodies against ovalbumin were detected by enzyme-linked immunosorbent assay. In the positive control, Freund's adjuvant was used.

Statistics

Student's *t* test was used to statistically analyze the data.

RESULTS

As can be observed in the elemental analysis and particularly in the ¹³C-NMR analysis, HZ presents more functionality than HC in both the aromatic and aliphatic moieties. This functionality is expressed in higher contents of phenolic carbon, glycol carbon, and alkyl carbons. However, HC presents a marked aromatic character, whereas HZ presents a predominant aromatic character with significant aliphatic arrangements. Thus, we have two HA with very different structures. HZ is more functionalized and with combining aromatic and aliphatic moieties, and HC is less functionalized but has high aromatic character (Tables 1 and 2).

First, we measured the effects of HA with and without glucan on numbers of cells in the peritoneal cavity. Using the same combinations as on all subsequent experiments, we observed no changes in cellularity (with respect to both total numbers and differential counts) after either intraperitoneal or oral application (data not shown). In all cases, glucan, HA, or both were dissolved in PBS, which was also used as a negative control.

The effects of various glucans on macrophages are well established. However, in order to demonstrate that a new generation of immunomodulators really exhibits an immunomodulatory property, an evaluation of phagocytosis is necessary. First, we measured the effects of glucan and/or HA on *in vitro* phagocytosis of synthetic HEMA microspheres in peripheral blood (Fig. 1). Both glucan and HA stimulate the internalization of synthetic particles, but the combined preparation, particularly in the 1:1 ratio, exhibited a significant synergistic effect on blood neutrophils. Identical results were achieved when we measured *in vitro* phagocytosis of peritoneal macrophages (Fig. 2). In both cases, the cells were isolated from mice injected with glucan and/or HA.

Evidence of the immunomodulating activity was also demonstrated through effects on the production of IL-2 by spleen cells (Fig. 3) and the production of IL-2 was measured

TABLE 1. ELEMENTARY ANALYSIS OF HA SAMPLES

HA	%C	%H	%N	%O ^a
HZ	48.2	2.99	0.98	48.9
HC	57.5	1.60	1.10	39.8

^aBy difference.

TABLE 2. ¹³C-NMR SPECTROSCOPY OF HA SAMPLES

HA	Region (ppm)			
	Alkyl C-O-Alkyl (40-45)	Alkyl C (45-110)	Aromatic (110-160)	Carbonyl C (160-215)
HZ	27.4	12.6	46.5	12.8
HC	13.8	3.2	79.5	5.4




FIG. 1. Potentiation of *in vitro* phagocytosis of synthetic microspheres (HEMA particles) by intraperitoneally injected glucans and/or HA. A total of 100 μg of material in various ratios was injected. Control values (PBS only) were 33.6%. Peripheral blood neutrophils with three and more HEMA particles were considered positive. Data are mean ± SD values from three separate experiments (five mice per experimental group). *Significant differences at the *P* < .05 level.




FIG. 2. Potentiation of *in vitro* phagocytosis of synthetic microspheres (HEMA particles) by intraperitoneally injected glucans and/or HA. A total of 100 μg of material in various ratios was injected. Control values (PBS only) were 36.5%. Peritoneal macrophages with three and more HEMA particles were considered positive. Data are mean ± SD values from three separate experiments (five mice per experimental group). *Significant differences at the *P* < .05 level.




FIG. 3. Effects of glucan-HA combinations on Con A-stimulated secretion of IL-2 by spleen cells. The secretion of IL-2 by non-stimulated splenocytes (PBS group) is zero or very low. The lowest stimulation was statistically significant. Data are mean ± SD values from three separate experiments (five mice per experimental group).

TABLE 3. EFFECT OF GLUCAN-INDUCED CYTOKINE SECRETION

HA	Glucan	IL2	IL4	IL5	IL6	TNF- α	MCP-1
HZ							
100%	0%	111.2 \pm 8.8	149.2 \pm 1.8	77.1 \pm 4.4	56.2 \pm 2.3	109.1 \pm 5.5	1.0 \pm 0.1
90%	10%	233.8 \pm 14.9	155.6 \pm 8.8	56.6 \pm 2.2	50.5 \pm 4.5	120.1 \pm 6.7	8.1 \pm 1.1
50%	50%	534.6 \pm 21.2	651.5 \pm 21.8	268.8 \pm 11.5*	109.2 \pm 3.3*	434.4 \pm 21.7*	17.2 \pm 1.1*
10%	90%	599.9 \pm 44.5	601.2 \pm 33.8	301.6 \pm 22.0*	88.8 \pm 4.4*	443.4 \pm 19.9*	9.2 \pm 0.9*
0%	100%	597.2 \pm 44.8	580.9 \pm 40.1	119.6 \pm 8.1	48.2 \pm 3.8	270.1 \pm 9.9	6.0 \pm 1.1
HC							
100%	0%	36.6 \pm 1.8	97.1 \pm 4.4	21.2 \pm 1.9	61.5 \pm 3.5	76.5 \pm 3.9	0
90%	10%	148.4 \pm 8.8	142.2 \pm 5.5	26.1 \pm 2.2	38.4 \pm 4.1	134.4 \pm 7.1	1.0 \pm 0.1
50%	50%	465.4 \pm 22.3	712.5 \pm 34.9	199.7 \pm 9.1	101.1 \pm 4.5	354.5 \pm 18.7	9.1 \pm 1.7*
10%	90%	715.2 \pm 42.9	608.2 \pm 36.5	215.3 \pm 11.1*	188.4 \pm 9.9*	489.6 \pm 17.9*	4.4 \pm 0.8

Data are mean \pm SD values from three separate experiments (five mice per experiment) (PBS only) were always zero. *Represents significant differences between glucan-HA and either glucan only of HA-only groups at the $P \leq .05$ level.

after a 72-hour *in vitro* incubation of spleen cells isolated from control rats treated with or without a combination of glucan and HA resulted in significant stimulation of sIL-2 production. In the case of HZ, the highest stimulation was found with a 1:1 HZ:glucan ratio. In the case of HC, the highest production of IL-2 was found in the 1:9 ratio. All combinations showed much stronger stimulation than the HA or glucan alone. As the secretion of IL-2 by nonstimulated splenocytes (PBS group) is zero, even the lowest stimulation was statistically significant.

After the initial experiments, we measured the secretion of cytokines from mice injected intraperitoneally with the test material 24 hours earlier. The results shown in Table 3 clearly demonstrate a general important observation: both HA and glucan stimulated the secretion of only six of the 14 tested cytokines (IL-2, IL-4, IL-5, IL-6, IL-1, TNF- α , and MCP-1) with the only exception being HC, which did not stimulate production of MCP-1. With respect to HZ, combination with glucan resulted in higher stimulation of IL-5, IL-6, and TNF- α secretion than in the case of HZ or glucan alone. A highly different situation was found with the HC-glucan combination for which elevated levels of IL-5, IL-6, and TNF- α were found only in the 1:9 ratio. In both cases we found only a small stimulation of MCP-1 secretion.

We then focused on the use of our combinations as an adjuvant. As an experimental model, we used immunization with ovalbumin. Glucan, HA, or both were applied together with two intraperitoneal doses of antigen, the so-called Freund's adjuvant was used as an additional positive control. The results (Fig. 4) showed that the tested combination exhibited in all cases a significant adjuvant activity against antigen alone (optical density, 0.36 \pm 0.026). In the case of HZ, the highest stimulation was achieved with the 9:1 ratio, whereas in the case of HC, we found the highest stimulation with the 1:1 Freund's ratio (optical density, 1.67 \pm 0.22).

In the final step, mice challenged with D264 mammary tumors were tested for a therapeutic response (daily intraperitoneal injections of the tested substances (Fig. 5).

This experiment was repeated three times (three mice per experimental group) with similar results. The control values (PBS only) showed mean tumor weight of 699.7 \pm 38.5 mg. Our data showed that in the case of HZ, the molecule responsible for the strong inhibition of tumor growth is clearly the glucan, as glucan-caused inhibition was the same as with HZ:glucan combination. Regarding the HC:glucan combination, the latter showed the strongest inhibition of tumor growth from all tested samples. In both cases, HA alone showed no significant activity.

DISCUSSION

Humic substances are the main components of humus in the soil. They are produced by chemical and microbial degradation of organic matter coming from plants and ani-




FIG. 4. Effects of two intraperitoneal injections of tested HA-glucan combinations on the formation of antibodies against ovalbumin. Mice were injected twice (2 times separate) and serum was collected 7 days after last injection. The level of specific antibodies against ovalbumin was detected by enzyme-linked immunosorbent assay. As the positive control, Freund's adjuvant was used. Data are mean \pm SD values from three separate experiments (five mice per experimental group). *Significant differences between control (ovalbumin alone) and samples at the $P \leq .05$ level. OD, optical density.




FIGURE 5B. 5HAs glucan therapy of B16A/B7c tumor with H-Plat60a mammary carcinoma. Data from these independent experiments are shown (three groups per experimental group) for each treatment group of treated with each form of glucan as a therapy as indicated by the weight of tumors after 2 weeks of therapy. In each experiment, individual groups were given daily intraperitoneal injections of 100 mg of HA, glucan, or the combination of PHS and a group of mice given daily intraperitoneal injections of PBS had a tumor weight of 600 mg. Data are shown as the mean \pm SD values. Significant differences between treated and untreated groups at the P < .05 level.

...function of their solubility as a function of pH, three main organic humic fractions can be differentiated: HA which is soluble at alkaline pH but precipitates at acidic pH; fulvic acid which is soluble at any pH and humic, which is insoluble at any pH. The structure of these substances is not clear. From a qualitative point of view, they possess the simplest of functional groups and structural arrangements. However, in the polydispersed nature of the systems makes it difficult to suggest definite structures. In general, they are acidic polyelectrolytes with different molecular weights and sizes and with diverse chemical character. A special feature of these substances is their ability to affect the metabolic development of different organisms. Thus, effects of humic substances on plant metabolism, growth, activity, and animal development have been described in many other studies. However, studies relating humic substances to biological activity are scarce. Different studies have shown many practical ways to characterize humic substances is the elemental analysis of ^{13}C NMR and elemental analysis. We have applied these techniques to the study of HA and the other experiments. As can be observed in the elemental analysis and ^{13}C NMR spectra, HA presents a high aromaticity of the HC in both the aromatic and aliphatic regions. This functional activity is expressed in higher content of phenolic, aromatic carbon, and alkylate. However, HC presents a high aromatic character whereas HZ, presents a predominant aromatic character with significant aliphatic arrangements. Thus, we have used HA with a very different structure HZ as in 0-functionalized and with combining aromatic and aliphatic moieties, and HC is less functionalized but with high aromatic character (Tables 1 and 2).

Various types of immunomodulators exist in particular, and the known immunostimulatory phagocytosis. Therefore, the evaluation of this basic type of immunoreaction is important for the development of the effectiveness of any biologically active immunomodulator. We tested peripheral blood leukocytes and peritoneal macrophages for changes in phagocytosis in the peritoneal cavity based on HEM Au. These particles have a slight negative charge and a specific local specificity to the cell surface, while glutaraldehyde is a very effective phagocytosis inhibitor. The results of the present study have been verified by phase contrast electron microscopy. The results showed that the combined use of both substances has caused significant increases in phagocytosis of macrophages. The combined preparations showed a significant synergistic effect on both macrophages and neutrophils. The data show that the effects of a single injection of test substances, although additional experiments found that a similar effect can be observed after oral administration (data not shown). Studies performed in these experiments do not depend on direct cell activation but on the number and percentage of cell types in the peritoneal cavity did not change.

In addition to the direct effect of various cells of the immune system, the immunostimulatory action of β -glucans and other immune modulators is caused by potentiation of synthesis and release of several cytokines. First data focused on the stimulation of IL-2 production by spleen cells *in vitro* and found that HA-glucan combinations stimulated not only higher release of IL-2 than the tested substances alone but in the case of 90% HZ, with glucan release was even higher than with HA.

Data were evaluated the effects of cytokines of 14 different cytokines in serum. Depending on the HC:glucan ratio the stimulation HZ was more pronounced with HZ than with HC. However, in all cases the production of cytokines was higher than with HA alone. The most stimulated cytokines were IL-5, IL-6, and IL-10; eight of the total of 14 tested cytokines were not detected.

Cytokines are important intercellular communicators, and their crucial role in immune response between different parts of immune system is well established. Reports on the role of IL-2 in tumor growth inhibition and growth promoting properties. The literature of a high rate of progression in experimental, and cytokines produced by malignant cells can function as both a growth factor and an immunomodulator.

Our results have established that glucan also supports the growth of tumor cells. In this immunoreaction by serum cytokines, we compared the adjuvant effect of HA and the tested glucan-HA combinations with that of Freund's adjuvant. Our results show that although the activities were always lower than those of Freund's adjuvant, they were more effective in significant with the highest activity found with the combination of glucan-HC.

Finally, we decided to test, if possible, effects of HA-glucan combinations *in vivo* growth of the mouse breast tumor cells. Our previous work has demonstrated that there is a high similarity of mouse and human TR3 in

response to glioxans, which makes the mouse tumor models suitable for investigation of glioxans.³⁰ We used the same experimental design as published before using yeast-derived glioxan.³¹ Similar to the antibody response, the highest effect was found with 20 mg/kg of glioxan, where we observed a 92% inhibition of tumor growth. This result indicates that aromaticity could play an important role in this antitumor effect of HAAs because HZ, which presents a significantly lower aromaticity than HC, did not show this activity.

Lipopolysaccharide contamination might mask the real effects of any immunomodulator. Therefore, we checked the lipopolysaccharide contamination of our solutions. Observed values were on a lower than 0.1 ng/ml. In addition, we functionally depleted lipopolysaccharide from both glioxan and HA by treatment with 10 µg/ml polymyxin B. We found identical results in all cases. The similarity between results obtained with regular and lipopolysaccharide-free material indicated that the minor lipopolysaccharide presence is not responsible for elevation of immunological activities and/or the antitumor response.

To summarize our data, our report suggests that HAAs are biologically active immunomodulators affecting both the humoral and cellular branches of immune reactions. In addition, both HZ and HC are working in synergy in stimulation of the immune reaction, a finding supporting further studies of these natural immunomodulators.

AUTHOR DISCLOSURE STATEMENT

No competing financial interests exist.

REFERENCES

1. Bonacera F, Sebastyane MM. Effect of bacterial endotoxins on the reticuloendothelial system. *Fed Proc* 1957; 16: 860–867.
2. Rigi SJ, Di Nuzio NR. Identification of a urothelial cell stimulating agent in pyrazosin. *Ann N Y Acad Sci* 1961; 200: 297–300.
3. Di Luzio NR, Williams DJ, McNamara BB, Edwards BF, Kitahama AI. Glucoparatic tumor-inhibitory and anti-bacterial activity of soluble and particulate glucan. *Int J Cancer* 1979; 24: 773–779.
4. Mimura H, Ohtsuki S, Suzuki Y, Adamae E. Purification, antitumor activity, and structural characterization of β -1,3- β -glucan from *Parakeet vesiculosa*. *Chem Pharm Bull* 1985; 33: 5096–5099.
5. Novak M, Vetvička V. Glucans as biological response modifiers. *Endocr Metab Immune Disord* 2009; 9: 637–75.
6. Dittova C, Bečejny S, Mrcková G. Modulation of diver fibrosis and pathological changes in mice infected with *Mycobacterium avium* after administration of glioxan and liposaccharide glucan in combination with vitamin C. *J Helminthol* 2003; 77: 219–226.
7. Velebný S, Tomášková V, Holkové A, Dubenský B, Terecárková I. Lipomycin and liposaccharide as immunomodulators: able to enhance the larvicidal effect of the anthelmintic? *Helminthologia* 1997; 34: 147–153.
8. Kukula M, Szatmari Z, Lutterová M, Kubová J, Vajdová K, Horáková E. Effects of floszofragin on endotoxin-enhanced gold ischemia reperfusion injury of rat liver. *Physiol Res* 2004; 53: 431–437.
9. Bittner M, Janáček J, Hölcková I, Čížek J, Holubek J, Ptáček L. Activation of TLR4 receptor by pure trimucic acids. *Environ Toxicol* 2006; 21: 338–342.
10. Bernacchi F, Bonzanelli M, Minuzzi M, Falezza A, Loprieno N, Barale R. Effect of acetylated glioxan on the natural humic acid. *Mutagenesis* 1996; 11: 467–469.
11. Helsen G, Schoneveld CHM, Le Chen J, Janssen M, Kalka G. Humic acid induces genotoxicity in peripheral blood lymphocytes using comet and sister chromatid exchange assay. *J Hazard Mat* 2008; 153: 784–792.
12. Gao J, Yang H, Sun S, Chen S, Sun J, Li B, Fu K. Humic acid suppresses the IL-1 β -induced expression of cell surface adhesion proteins through the inhibition of NF- κ B activation. *Biocol Appl Pharmacol* 2000; 166: 59–67.
13. Madej JA, Kuryśka J, Garbaliński T. The influence of long-term administration of tolpaepat preparation on immune reactivity in mice. *Acta Morphologica et Anatomica Polonica* 1993; 50: 397–404.
14. Chen C, Liu J, Yan J, Yang M, Lee Y, Huang T. The effect of humic acid on the adhesibility of neutrophils. *Thromb Res* 2003; 108: 67–76.
15. Kleckings P, Spang M, Witaler P, Eljeh K, Hildebrandt K. Antiviral activity of humic substances. *Z Physiother* 1983; 33: 95–101.
16. Joons G, Dekkers J, Jansen van Rensburg G. Investigation of the immunostimulatory properties of zoehumate. *Z Naturforsch C* 2003; 58: 263–267.
17. Islam KM, Schumacher O, Groppe J. Humic acid substances in animal agriculture. *Food Sci Nutr* 2005; 4: 126–134.
18. Vetvička V, Votava S, Sotava O, Šiška A, Váňová A, Váňová Z, Váňová Z, Křížová J, and resveratrol complectible possible synergistic effects on the immune system. *Biomed Pap Med Fac* 2007; 151: 41–46.
19. Vetvička V, Farnusová K, Šiška A, Kamáková J, Kasperk L, Mrazová M. Phagocytosis of human blood leukocytes: a simple method. *Immunol Lett* 1982; 5: 97–100.
20. Vetvička V, Mlýnský M, Křivánek H, Šiška A, Křivánek R. Alpha-fetoprotein and phagocytosis in thymic mice. *Immunol Lett* 1988; 19: 95–98.
21. Stevenson H. *Humus Chemistry: Composition, Reactions, and Properties*. Wiley-Interscience, New York, 1982.
22. Chen A. A review of the effects of humic substances on plant growth. In: *Humic Substances in the Environment*. Springer Science & Technology, American Society of Agronomy, Madison, WI, 1990, pp. 161–186.
23. Piccolo HM. Humus and soil: considerations of the humus substances in the soil. *Soil Science Society of America* 1996, pp. 225–264.
24. Sessé N, Miano F, Murek S. *Substance and the Global Environment and Implications on Human Health*. Elsevier, Amsterdam, 1994.
25. Aebel G, Sotava S, Chihara G, Frachet E. Effect of lentipato and mannitol on phagocytosis of fluorescent latex particles by mouse peritoneal macrophages. *Int J Immunopharmacol* 1989; 11: 615–621.
26. Tromba M, Ohashi K, Shimizu N, Gonda R. Characterization of a novel glioxan which exhibits reticuloendothelial system-potentiating and anti-complementary activities from the azobite of *Cydonia oblongifolia*. *Chem Pharm Bull (Tokyo)* 1994; 42: 630–633.
27. Vetvička V, Farnusová K, Šiška A, Křivánek R. Biomedical applications of humic acid. *Biomaterials* 1987; 8: 343–345.

